This Page Is Inserted by IFW Operations and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.

			•
A6.1K	9/00MC20B	/5-1 K	9/120
€09K-	-3/30	EΡ	0588897
WO	9222288	AUD	2017592
CA	2111002	СŅ	1067578
C.Z	9302714	FIDO	935464
NO	934500		

しつようつじゅ



11 Publication numb r: 0 518 600 A1

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 92305252.6

(51) Int. CI.5: A61K 9/72, C09K 3/30

(22) Date of filing: 08.06.92

(30) Priority: 10.06.91 US 712791

43 Date of publication of application: 16.12.92 Bulletin 92/51

84 Designated Contracting States :

7) Applicant: SCHERING CORPORATION 2000 Galloping Hill Road Kenilworth New Jersey 07033 (US)

72 Inventor: Fassberg, Julianne
175W. 12th Street
New York, N.Y. 10011 (US)
Inventor: Sequeira, Joel A.
18 Stuyvesant Oval
New York, N.Y. 10009 (US)
Inventor: Chaudry, Imtiaz A.
18 Rose Avenue
North Caldwell, New Jersey 07006 (US)
Inventor: Kopcha, Michael
141 Wycoff Way West
East Brunswick, New Jersey 08816 (US)

(4) Representative: Ritter, Stephen David et al Mathys & Squire 10 Fleet Street London EC4Y 1AY (GB)

- (54) Non-chlorofluorocarbon aerosol formulations.
- (57) Aerosol formulations substantially free of chlorofluorocarbons for oral and/or nasal administration are described. The formulations comprise 1,1,1,2,3,3,3 heptafluoropropane, a medicament, optionally an excipient and optionally a surfactant. Methods of treatment utilizing the formulations also are described.

EP 0 518 600 A1

INTRODUCTION TO THE INVENTION

10

15

20

30

50

55

The present invention is directed at aerosol formulations which are substantially free of chlorofluorocarbons (CFC's). More specifically, the present invention is directed at formulations substantially free of CFC's and having particular utility in medicinal applications, especially in metered dose pressurized inhalators (MDI's).

Metered dose inhalators have proven to be an effective method for delivering medicaments orally and nasally. They have been used extensively for delivering bronchodilating and steroidal compounds to asthmatics and may also be useful for delivering other compounds such as pentamidine and non-bronchodilator anti-inflammatory drugs. The rapid onset of activity of compounds administered in this manner and the absence of any significant side effects have resulted in a large number of compounds being formulated for administration via this route. Typically, the drug is delivered to the patient by a propellant system generally comprising one or more propellants which have the appropriate vapor pressure and which are suitable for oral or nasal administration. The more preferred propellant systems typically comprise propellant 11, propellant 12, propellant 114 or mixtures thereof. Often the vapor pressure of the propellant systems is adjusted by admixing a liquid excipient with the propellant.

However, propellants 11, 12 and 114 belong to a class of compounds known as chlorofluorocarbons, which have been linked to the depletion of ozone in the atmosphere. It has been postulated that ozone blocks certain harmful UV rays and that a decrease in the atmospheric ozone content will result in an increase in the incidence of skin cancer. In the 1970's certain steps were taken to reduce the CFC emissions from aerosols. Other propellants, such as hydrocarbons, were used, or the product was delivered in a different manner. Because CFC usage in medicinal applications is relatively low i.e. less than 1% of total CFC emissions, and because of the health benefits associated with metered dose inhalators, steps were not taken at that time to restrict the use of CFC propellants in metered dose inhalators.

However, continuing and more sophisticated ozone measurements have indicated that the earlier restrictions in CFC usage were insufficient and that additional, significant steps should be taken to drastically reduce CFC emissions. Recently, recommendations have been made that CFC production be virtually discontinued by the end of this century. As a result, it may not be possible to continue to use CFC propellants in the intermediate and long term. While some efforts have been made to use non-pressurized metered dose inhalators, many of these devices have not been completely successful. Many do not deliver uniform doses, are mechanically complex, do not provide the 100-200 doses per unit of current aerosol containers, are difficult for individuals to utilize, and are bulky and/or cumbersome for the patients to use, particularly when they have an acute need for the medication.

As a result, there is a need for aerosol formulations which are substantially free of CFC's. Non-CFC propellants systems must meet several criteria for pressurized metered dose inhalators. They must be non-toxic, stable and non-reactive with the medicament and the other major components in the valve/actuator. One propellant which has been found to be suitable is CF₂-CH₂F-CF₃, also known as Freon 227, HFA 227, HFC 227 or 1,1,1,2,3,3,3 heptafluoropropane. However, certain physical properties, i.e., polarity and solubility, of HFC 227 differ from those of commonly used CFC propellants. Commonly used surfactants may be insoluble in HFA 227. Moreover, where the medicament is to be delivered as a solution, the medicament may not be readily soluble in this propellant. The polarity difference between HFC 227 and the previously used CFC propellants may result in a different delivery of the medicament when HFC 227 replaces a CFC propellant. The medicament may cream, settle or agglomerate in the non-CFC propellant even though this did not occur in the CFC propellant.

The use of HFA 227 previously has been disclosed for use in medicinal inhalators. European Patent Publication No. 0 384 371 is directed at the combination of propellant 227 and propane, butane, isobutane, Me_2O and/or F_2CHMe .

Research Disclosure No. 30161, May, 1989 discloses that non-CFC propellants, such as fluorohydrocarbons may be used in pressurized medicaments delivered directly to the lungs, e.g. bronchodilators.

Other publications have been directed at the use of other fluorohydrocarbons, such as HFC 134a, for aerosol propellants. European Patent Publication No. 0 372 777 is directed at medicinal aerosol formulations incorporating HFC 134a and an adjuvant having a higher polarity than the propellant. This publication lists several possible adjuvants and surfactants for use in combination with the propellant and the medicament.

International patent application No. WO 91/04011 discloses the combination of HFC 134a and a powdered medicament pre-coated with a non-perfluorinated surfactant prior to dispersing the powdered medicament in the propellant. Pages 6-7 of the publication list suitable surfactants for use with the propellant. A perfluorinated adjuvant optionally could be added. How v r, the pre-coating of the medicament may not be advantageous, since it adds an additional, complex step to the manufacturing process.

U.S. Patent No. 4,174,295 discloses the combination of HFC 134a with various chlorofluorocarbons and

optionally a saturated hydrocarbon. U.S. Patent No. 2,885,427 discloses the use of HFC-134a as an aerosol propellant. U.S. Patent No. 3,261,748 discloses the use of HFC-134a for anesthesia. U.S. Patent Nos. 4,129,603, 4,311,863, 4,851,595 and European Publication No. 379,793 also disclose the use of HFC-134a as an aerosol propellant.

However, the specific combinations noted above may not provide the desired solubility, stability, low toxicity, exact dosage, correct particle size (if suspension) and/or compatibility with commonly used valves assemblies of metered dose inhalers.

SUMMARY OF THE INVENTION

10

15

20

25.

30

35

40

50

55

5

Accordingly, the present invention is directed at a non-toxic formulation substantially free of CFC's having improved stability and compatibility with the medicament and which is relatively easily manufactured.

The present invention also is directed at formulations which may be utilized in present aerosol filling equipment with only relatively minor modifications and without pre-coating the medicament.

One embodiment of the present invention is directed at a formulation comprising:

A. Propeliant 1,1,1,2,3,3,3 heptafluoropropane;

B. optionally an excipient selected from the group consisting of alcohols, Miglyol 812, Miglyol 840, PEG-400, menthol, lauroglycol, Vertrel 245. Transcutol, Labrafac Hydro WL 1219, perfluorocyclobutane, eucallyptus oil, short chain fatty acids, and combinations tereof;

C. a medicament, and

D. optionally a surfactant selected from the group consisting of oleic acid, sorbitan trioleate, cetyl pyridinium chloride, soya lecithin, Tween 20, Tween 60, Tween 80, Pluronic L-121 and Pluronic L-92, castor oil ethoxylate, pluronic F 68, Tetronic 150 R1 and combinations thereof.

Also included within the invention is an aerosol formulation comprising:

A. an effective amount of medicament,

B. 1,1,1,2,3,3,3 heptafluoropropane; and

C. an excipient selected from the group consisting of: propylene glycol diesters of medium chain fatty acids;

triglyceride esters of medium chain fatty acids;

perfluorodimethylcyclobutane;

perfluorocyclobutane;

polyethylene glycol;

menthol;

lauroglycol;

diethylene glycol monoethylether;

polyglycolized glycerides of medium chain fatty acids;

alcohols:

eucalyptus oil;

short chain fatty acids;

and combinations thereof.

The formulation optionally may further comprise a surfactant. The surfactant preferably is selected from the group consisting of:

oleic acid;

sorbitan trioleate;

cetyl pyridinium chloride;

45 soya lecithin;

polyoxyethylene(20) sorbitan monolaurate;

polyoxyethylene (10) stearyl ether,

polyoxyethylene (2) oleyl ether,

polyoxypropylene-polyoxyethylene-ethylene diamine block copolymers;

polyoxyethylene(20) sorbitan monostearate;

polyoxyethylene(20) sorbitan monooleate;

polyoxypropylene-polyoxyethylene block copolymers;

castor oil ethoxylate; and combinations thereof.

The preferred liquid excipients are diethylene glycol monethyether, propyleneglycol diesters of medium chain fatty acids, perfluorodimethylcyclobutane and polyethylene glycol.

The preferred surfactants are oleic acid; sorbitan trioleate, cetylpyridinium chloride; polyoxyethylene (20) sorbitan monolaurate; polyoxypropylene-polyoxyethylene block copolymers; soya lecithin; and polyoxypropylene-polyoxyethylene-ethylenediamine block copolymers; with oleic acid being particularly preferred.

The invention is of particular utility where the medicament is albuterol, mometasone furoate or beclome-thasone dipropionate, and salts and clathrates thereof.

A useful formulation range comprises:

A. 1,1,1,2,3,3,3 heptafluoropropane 25 - 99.99 wt %

B. medicament 0.01 - 1 wt %

C. excipient 0 - 75 wt %

D. surfactant 0 - 3 wt %

The present invention also is directed at a method of treating asthma in mammals comprising administering to a mammal in need of such treatment an effective amount of aerosol formulation comprising:

A. a medicament selected from the group comprising albuterol, mometasone furoate, beclomethasone dipropionate, and salts and dathrates thereof;

B. 1,1,1,2,3,3,3 heptafluoropropane; and

C. optionally an excipient selected from the group consisting of:

propylene glycol diesters of medium chain fatty acids;

triglyceride esters of medium chain fatty acids;

perfluorodimethylcyclobutane;

perfluorocydobutane;

polyethylene glycol;

menthol;

10

15

20

25

30

35

40

45

50

lauroglycol;

diethyleneglycol monoethylether,

polyglycolized glycerides of medium chain fatty acids;

alcohols:

short chain fatty acids;

eucalyptus oil; and combinations thereof.

A surfactant optionally is present. The surfactant preferably is selected from the group consisting of:

oleic acid;

sorbitan trioleate;

cetyl pyridinium chloride;

soya lecithin;

polyoxyethylene (20) sorbitan monolaurate;

polyoxyethylene (10) stearyl ether;

polyoxyethylene (2) oleyl ether,

polyoxyethylene-polyoxypropylene-ethylene diamine block copolymers;

polyoxyethylene (20) sorbitan monostearate;

polyoxypropylene-polyoxyethylene block copolymers;

castor oil ethoxylate; and combinations thereof.

DETAILED DESCRIPTION OF THE INVENTION

The formulations of the present invention all utilize propellant 227 in combination with the medicament, optionally a liquid excipient and optionally a surfactant.

The excipient facilitates the compatibility of the medicament with the propellant and also lowers the discharge pressure to an acceptable range i.e. about 2.76 - 5.52 x 105 newton/meter² absolute (40 to 80 psia), preferably 3.45 - 4.83 x 105 newton/meter² absolute (50 to 70 psia). The excipient chosen must be non-reactive with the medicament, relatively non-toxic, and should have a vapor pressure below about 3.45 x 105 newton/meter² absolute (50 psia). As used hereinafter the term "medium chain fatty acids" refers to chains of alkyl groups terminating in a -COOH group and having 6-12 carbon atoms, preferably 8-10 carbon atoms. The term "short chain fatty acids" refers to chains of alkyl groups terminating in a -COOH group and having 4-8 carb in atoms. The term "alcohol" includes C₁-C₃ alcohols, such as methanol, ethanol and isopropanol. Among the preferred excipients are:

propylene glycol diesters of medium chain fatty acids available under the tradenam Miglyol 840 (from Hūls America, Inc. Piscataway, N.J.);

triglyceride esters of medium chain fatty acids available under the tradename Miglyol 812 (from Hūls);

perfluorodimethylcyclobutane available under the tradename Vertrel 245 (from E.I DuPont de Nemours and Co. Inc. Wilmington, Delaware);

perfluorocyclobutane available under the tradename octafluorocyclobutane (from PCR Gainsville, Florida);

polyethylene glycol available under the tradename PEG 400 (from BASF Parsippany, N.J.); menthol (from Pluess-Stauffer International Stanford, Connecticut);

propylene glycol monolaurate available under the tradename lauroglycol (from Gattefossé Elmsford, N.Y.);

diethylene glycol monoethylether available under the tradename Transcutol (from Gattefossé); polyglycolized glyceride of medium chain fatty acids available under the tradename Labrafac Hydro WL 1219 (from Gattefossé);

alcohols, such as ethanol, methanol and isopropanol;

5

10

15

20

25

30

35

40

45

50

55

eucalyptus oil available (from Pluess-Stauffer International); and mixtures thereof.

A surfactant optionally may be added to lower the surface and interfacial tension between the medicament and the propellant. Where the medicament, propellant and excipient are to form a suspension, a surfactant may or may not be required. Where the medicament, propellant and excipient are to form a solution, a surfactant may or may not be necessary, depending in part, on the solubility of the particular medicament and excipient. The surfactant may be any suitable, non-toxic compound which is non-reactive with the medicament and which substantially reduces the surface tension between the medicament, the excipient and the propellant and/or acts as a valve lubricant. Among the preferred surfactants are:

oleic acid available under the tradename oleic acid NF6321 (from Henkel Corp. Emery Group, Cincinnati, Ohio);

cetylpyridinium chloride (from Arrow Chemical, Inc. Westwood, N.J.);

soya lecithin available under the tradename Epikuron 200 (from Lucas Meyer Decatur, Illinois);

polyoxyethylene(20) sorbitan monolaurate available under the tradename Tween 20 (from ICI Specialty Chemicals, Wilmington, Delaware);

polyoxyethylene(20) sorbitan monostearate available under the tradename Tween 60 (from ICI); polyoxyethylene(20) sorbitan monooleate available under the tradename Tween 80 (from ICI); polyoxyethylene (10) stearyl ether available under the tradename Brij 76 (from ICI);

polyoxyethylene (2) oleyl ether available under the tradename Brij 92 (from ICI);

polyoxyethylene-polyoxypropylene-ethylenediamine block copolymer available under the tradename Tetronic 150 R1 (from BASF);

polyoxypropylene-polyoxyethylene block copolymers available under the tradenames Pluronic L-92, Pluronic L-121 and Pluronic F 68 (from BASF);

castor oil ethoxylate available under the tradename Alkasurf CO-40 (from Rhone-Poulenc Mississauga Ontario, Canada); and mixtures thereof.

The medicaments of the present invention may include any pharmaceutically active compounds which are to be delivered by oral inhalation or nasally. Typical classes of compounds include bronchodilators, anti-inflammatory compounds, antihistamines, antiallergics, analgesics, antitussives, anti-anginal medications, steroids, corticosteroids, vasoconstrictors and antibiotics. Specific compounds within these classes of compounds are albuterol, mometasone furoate, beclomethasone dipropionate, isoproterenol, heparin, terbutaline, rimiterol, perbuterol, disodium cromoglycate, isoprenaline, adrenaline, pentamidine and ipratropium bromide. These compounds may be utilized either as the free base, as a salt, or as a clathrate, depending upon the stability and solubility of the active compound in the specific formulation. When clathrates are utilized, P-11 and hexane clathrates are particularly preferred.

Where the active compound forms a suspension, the particle size should be relatively uniform, with substantially all the particles preferably ranging between about 0.1-25 microns, preferably 0.5-10 microns, more preferably 1-5 microns. Particles larger than 25 microns may be held up in the oropharyngeal cavity, while particles smaller than about 0.5 micron preferably are not utilized, since they would be more likely to be exhaled and, therefore, not reach the lungs of the patient.

The formulations of the present invention may be filled into the aerosol containers using conventional filling equipment. Since propellant 227 may not be compatible with all elastomeric compounds currently utilized in present aerosol valve assemblies, it may be necessary to substitute other materials, such as white buna rubber, or to utilize excipients and optionally surfactants which mitigate the adverse effects of propellant 227 on the valve components.

To assure uniform dispersion of the active ingredient, the formulations typically will include the following components:

_		Range (wt %)	Preferred Range (wt%)	Most Preferred Range (wt%)
5	Medicament Propellant	0.01 - 1 25 - 99.99	0.03 - 0.7 50 - 99.97	0.05 - 0.5 50 - 99.95
10	Excipient(s) Surfactant(s)	0 - 75 0 - 3	0 - 50 0 - 2	0 - 50 0 - 1

Depending on the particular application, the container may be charged with a predetermined quantity of formulation for single or multiple dosing. Typically, the container is sized for multiple-dosing, and, therefore, it is very important that the formulation delivered is substantially uniform for each dosing. For example, where the formulation is for bronchodilation, the container typically is charged with a sufficient quantity of the formulation for 200 charges.

Suitable suspensions may be screened in part by observing several physical properties of the formulation, i.e. the rate of particle agglomeration, the size of the agglomerates and the rate of particulate creaming/settling and comparing these to an acceptable standard. Suitable solutions may be screened by observing the solubility of the medicament over the entire recommended storage temperature range.

Suspensions of the present invention preferably may be prepared by either the pressure filling or cold filling procedures well-known in the art.

For metered dose inhalators, suspensions may be particularly preferred for efficacy and stability considerations.

Those skilled in the art may choose to add one or more preservative, buffer, antioxidant, sweetener and/or flavors or other taste masking agents depending upon the characteristics of the formulation.

Examples I - XXXIII below further describe the present invention. For several of the examples, alternative formulations denoted as A and B are provided.

Component

₩

35

EXAMPLE I

40		A	B	
	Albuteroi	0.5	0.1	
	Miglyol 812	10.0	1.0	
	HFC-227	89.5	98.9	
45	- •	10.0	1.0	

EXAMPLE II

50	Albuterol	0.1
	Transcutol	25.0
	HFC-227	74.9

55

25

EXAMPLE III

5	•		A	В
	Albuterol		0.5	0.1
	Miglyol 840		10.0	1.0
	HFC-227		89.5	98.9
10				
		EXAMPLE IV		
15	•			
	Albuterol		0.1	
	PEG 400		1.0	
	HFC-227		98.9	
20				
		EXAMPLE V		
25 .	Albuterol		0.1	
	Menthol		0.5	
	HFC 227		98.9	
<i>30</i>				
		EXAMPLE VI		
			Α	В
35	Albuterol		0.1	0.1
	Lauroglycol		0.1	0.5
	HFC 227		99.8	99.4
40				
		EXAMPLE VII		
45			Α	B
	Albuterol		0.1	0.5
	Vertrel 245		10.0	49.6
	HFC 227		89.9	49.9
50				

EXAMPLE VIII

5	Albuterol		0.1	
	Labrafac Hydro WL 1219		0.5	
	HFC 227		99.4	
10				
,,		EXAMPLE IX		
			Α	B
15	Albuterol		0.1	0.5
	Perfluorocyclobutane		10.0	49.6
	HFC 227		89.9	49.9
20		EXAMPLE X		
		CARMIFLE A		
· .			Α	В
25	Oleic Acid		0.01	0.1
	Albuterol		0.10	0.1
	Ethanol		1.00	30.0
30	HFC 227		98.89	69.8
		EXAMPLE XI		
35			A	B
	Oleic Acid		0.01	0.1
	Albuterol sulfate		0.10	0.1
	Ethanol		1.00	30.0
40	HFC 227		98.89	69.8
		EXAMPLE XII		
45			Α	B
	Oleic Acid		0.01	0.1
	Albuterol		0.10	0.1
	Ethanol		1.00	25.0
50	HFC 227		98.89	74.8

EXAMPLE XIII

5			Α	88
	Oleic Acid		0.001	0.01
	Albuterol		0.1	0.1
	Miglyol 812		1.0	10.0
10	HFC 227		98.8	89.8
		EXAMPLE XIV		
15				
	Tetronic 150 R1		0.1	
	Albuterol		0.1	
20	Miglyol 812		9.8	
•	HFC-227		90	
25		EVALADI E sa		
		EXAMPLE XV	•	_
	Pluronic L121		A	B
30	Albuteroi		0.1	0.1
30	Miglyol 812	·	0.1	0.1
	HFC 227		1.0	10.0
	• 12.	•	98.8	89.8
35		EXAMPLE XVI		
	Tween 20		0.1	
40	Albuterol		0.1	
	Miglyol 812		10.0	
	Vertrel 245		10.0	
45	HFC-227		79.8	

50

EXAMPLE XVII

		A	В
5	Oleic Acid	0.01	0.1
	Albuterol Sulfate	0.10	0.1
	Ethanol	1.00	25.0
10	HFC 227	98.89	74.8
.0		EXAMPLE XVIII	
		A	В
15	Oleic Acid	0.01	0.1
	Albuterol Sulfate	0.10	0.1
	Transcutol	1.00	25.0
20	HFC 227	98.89	74.8
		EXAMPLE XIX	
		A	<u> </u>
25	Pluronic L 121	0.1	0.1
	Mometasone Furoate	0.1	0.1
	Miglyol 812	1.0	10.0
30	HFC 227	98.8	89.8
		EXAMPLE XX	
35			
•	Tetronic 150 R1	0.1	
	Mometasone Furoate	0.1	
	Miglyol 812	9.8	
40	HFC-227	90	
		EXAMPLE XXI	
45			
	Mometasone Furoate	0.1	
	HFC-227	99.9	

55

EXAMPLE XXII

5	Beclomethasone Dipropionate HFC-227	0.1 99.9
10	EXAMPLE	≣ XXIII
15	Mometasone Furoate Tween 20 HFC-227	0.1 0.01 99.89 E XXIV
20	Beclomethasone Dipropionate Tween 20 HFC-227	0.1 0.01 99.89
25	EXAMPL	EXXV
30	Mometasone Furoate Tween 20 Oleic Acid HFC-227 EXAMPL	0.1 0.01 0.0005 99.8895 .E XXVI
35	Beclomethasone Dipropionate	0.1
40	Tween 20 Oleic Acid HFC-227	0.01 0.0005 99.8895
45	EXAMPLE	EXXVII
50	Mometasone Furoate Miglyol 812 Oleic Acid Tetronic 150 R1 HFC-227	0.1 9 0.005 0.01 90.885

EXAMPLE XXVIII

5	Beclomethasone Dipropionate Miglyol 840 Oleic Acid Pluronic L121 HFC-227	0.1 9 0.005 0.01 90.885	
	EXAMP	LE XXIX	
15		A	В
	Oleic Acid	0.001	0.01
	Mometasone Furoate	0.1	0.1
20	Miglyol 812 HFC 227	1.0	10.0
	111 0 227	98.8	89.8
25	EXAMP	LE XXX	
	D	A	B
	Pluronic L121	0.1	0.1
30	Beclomethasone Dipropionate	0.1	0.1
	Miglyol 812 HFC 227	1.0	10.0
	1110 227	98.8	89.8
35	EXAMPL	E XXXI	
		A	В
40	Beclomethasone Dipropionate	0.1	0.1
	Miglyol 812	1.0	10.0
	HFC 227	98.9	89.9
45	EXAMPLE	EXXXII	
	_	Α	В
50	Beclomethasone Dipropionate	0.1	0.1
	PEG 400	1.0	10.0
	HFC 227	98.9	89.9

EXAMPLE XXXIII

Beclomethasone Dipropionate	0.1
Ethanol	5
HFC 227	94.9

While the examples above have been directed at albuterol, albuterol sulfate, mometasone furoate, bedomethasone dipropionate and bedomethasone dipropionate dathrates, it is contemplated that other orally or nasally administered medicaments could be utilized. Similarly, it is contemplated that excipients and surfactants other than those exemplified may be utilized.

The descriptions of the foregoing embodiments of the invention have been presented for the purpose of illustration and description. They are not intended to be exhaustive or to limit the invention to the precise forms disclosed, and obviously many modifications and variations are possible in light of the above teaching. The embodiments were chosen and described in order to best explain the principles of the invention and its practical application to thereby enable others skilled in the art to best utilize the invention in various embodiments and with various modifications as are suited to the particular use contemplated. It is intended that the scope of the invention be defined by the claims appended hereto.

Claims

5

15

20

35

40

50

55

25 1. An aerosol formulation consisting essentially of:

A. an effective amount of a medicament;

B. 1,1,1,2,3,3,3 heptafluoropropane; and optionally, one or more components selected from one or more of the following:

excipients;

30 surfactants; and

additives which are:

preservatives;

buffers:

antioxidants:

sweetners; and

taste masking agents.

2. The formulation of claim 1 wherein the excipient is selected from the group consisting of:

propylene glycol diesters of medium chain fatty acids;

triglyceride esters of medium chain fatty acids;

perfluorodimethylcyclobutane;

perfluorocyclobutane;

polyethylene glycol;

menthol;

45 lauroglycol;

diethylglycol monoethylether;

polyglycolized glycerides of medium chain fatty acids;

alcohols;

short chain fatty acids;

eucalyptus oil; and combinations thereof.

3. The formulation of claim 1 wherein the surfactant is selected from the group consisting of:

oleic acid;

s rbitan trioleate;

cetyl pyridinium chloride;

soya lecithin;

p ly xyethylene (20) sorbitan monolaurate;

polyoxyethylene(20) sorbitan monostearate;

polyoxyethylene(20) sorbitan monooleate; polyoxyethlyene (10) stearyl ether; polyoxyethylene (2) oleyl ether; polyoxyethylene-polyoxypropylene-ethylenediamine block copolymers; polyoxypropylene-polyoxyethylene block copolymers; castor oil ethoxylate; and combinations thereof.

- 4. The formulation of claim 1 wherein the medicament is selected from the group consisting of: albuterol, mometasone furoate, beclomethasone dipropionate, isoproterenol, heparin, terbutaline, rimiterol, perbuterol, disodium cromoglycate, isoprenaline, adrenaline, pentamidine, ipratropium bromide, and salts and clathrates thereof.
- 5. The formulation of claim 4 wherein the medicament is selected from the group consisting of:
 albuterol, albuterol sulfate, becomethasone dipropionate, becomethasone dipropionate clathrates
 and mometasone furoate.
 - 6. The formulation of claim 5 which is substantially free of chlorofluorocarbons.

5

10

15

35

45

- 7. The formulation of claim 5 containing an excipient selected from the group consisting of diethylene glycol monoethylether, propylene glycol diesters of medium chain fatty acids, perfluorodimethylcyclobutane and polyethylene glycol.
 - 8. The formulation of claim 7 containing a surfactant selected from the group consisting of: oleic acid, sorbitan trioleate, cetylpyridinium chloride and soya lecithin.
- 25 9. The formulation of claim 1 containing the following components in the indicated ranges:

	medicament	0.01-1 wt %
	1,1,1,2,3,3,3 heptafluoropropane	25 - 99.99 wt %
30	excipient	0 - 75 wt %
	surfactant	0 - 3 wt %

10. The formulation of claim 9 containing the following components in the indicated ranges:

	medicament	0.03 - 0.7 wt%
	1,1,1,2,3,3,3 heptafluoropropane	50 - 99.97 wt%
40	excipient	0 - 50 wt%
	surfactant	0 - 2 wt%

11. The formulation of claim 10 containing the following components in the indicated ranges:

	medicament	0.05 - 0.5 wt%
	1,1,1,2,3,3,3 heptafluoropropane	50 -99.95 wt%
50	excipient	0 - 50 wt%
	surfactant	0 - 1 wt%

- 12. The formulation of claim 9 wherein the medicam int is a powder having a mean particle size of about 1-5 microns.
 - 13. A method for treating mammals comprising administering to said mammals an effective amount the aerosol formulation of claim 1.

14. A m thod of treating asthma in mammals comprising administering to a mammal in need of such treatment an effective amount of aerosol formulation consisting essentially of: A. medicament selected from the group comprising albuterol, mometasone furoate, beclomethasone dipropionat, and salts and dathrates ther of, 5 B. 1,1,1,2,3,3,3 heptafluoropropane; C. optionally an excipi nt selected from th group consisting of: propylene glycol diesters of medium chain fatty acids; triglyceride esters of medium chain fatty acids; perfluorodimethylcyclobutane; 10 perfluorocyclobutane; polyethylene glycol; menthal: lauroglycol: diethylglycol monoethylether, 15 polyglycolized glycerides of medium chain fatty acids; alcohols: short chain fatty acids; eucalyptus oil; and combinations thereof: D. optionally a surfactant selected from the group consisting of: 20 oleic acid: sorbitan trioleate; cetyl pyridinium chloride; soya lecithin; polyoxyethylene (20) sorbitan monolaurate; . 25 polyoxyethylene (20) sorbitan monostearate: polyoxyethylene (20) sorbitan monooleate; polyoxyethylene (10) stearyl ether, polyoxyethylene (2) oleyl ether; polyoxyethylene-polyoxypropylene-ethylenediamine block copolymers; 30 polyoxypropylene-polyoxyethylene block copolymers; castor oil ethoxylate; and combinations thereof; and E. optionally one or more additives selected from at least one of the following classes: preservatives: buffers: antioxidants; 35 sweeteners; and taste masking agents. 40

15

45

50



PARTIAL EUROPEAN SEARCH REPORT

Application Number

which under Rule 45 of the European Patent Convention shall be considered, for the purposes of subsequent proceedings, as the European search report

EP 92 30 5252

Y: pa	uticularly relevant if combloed with an	after the fill other D : document of L : document o	ling date cited in the application ited for other reasons	
X: particularly relevant if taken alone after the Y: particularly relevant if combloed with another D: document document of the same category L: document			rinciple underlying the pit document, but publ	lavention ished on, or
TH	IE HAGUE	14-08-1992	BENZ	. K.F.
	Place of search	Date of completion of the second	•	Exempler
Remana a med body	rk: Although clai thod of treatment the search has b	ms 13,14 are dired of the human/animen carried out an effects of the con	mal nd	
The Sear the provi out a me Claims s Claims s	OMPLETE SEARCH rch Division considers that the present istons of the European Parent Convent maingful search into the state of the au earched completely: earched incompletely: for the limitation of the search:	European patent application does not co ion to such an extent that it is not possi it on the basis of some of the claims	omply with ble to carry	
				A 61 K C 07 K
	INT. GmbH) * Whole document, i example 5 *	n particular page 5, -/-	14	TECHNICAL FIELDS SEARCHED (Int. Cl.5)
Р,Х	* Page 1, line 1 - page 16, examples 8 WO-A-9 111 496 (BO	8-91; claims *	1-6,8-	
P,X	WO-A-9 114 422 (MI MANUFACTURING CO.)		1,4-6,9 -14	
P,X	WO-A-9 200 061 (FI * Page 1, line 1 - pages 8-10, example	page 8, line 2;	1-6,9-	
P,X	WO-A-9 200 062 (MI MANUFACTURING CO.) * Page 1, line 1 - page 11, example 47 examples 59-64; cla	page 6, line 39; ; pages 13,14,	1,2,4-6	A 61 K 9/72 C 09 K 3/30
ategory	Citation of document with in of relevant pas		Relevant to claim	CLASSIFICATION OF THE APPLICATION (Inc. CL.5)
		DERED TO BE RELEVA		

Off the William Co.

Page 2



PARTIAL EUROPEAN SEARCH REPORT

Apolication Number

EP 92 30 5252

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (lat. CL.5)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
Ρ,Χ	WO-A-9 111 495 (BOEHRINGER INGELHEIM INT. GmbH) * Whole document, in particular page 6, example 5 *	1-6,8-	
P,X	WO-A-9 111 173 (FISONS PLC) * Page 1, line 1 - page 9, line 10, pages 10,11, examples 2,4,6,8; claims *	1-14	
			TECHNICAL FIELDS SEARCHED (Int. CL5)